

Anti-Basal body marker [LhS 28]

Catalogue number: 151296

Sub-type: Primary antibody

Images:

Contributor

Inventor: Jenny Southgate ; Ludwik Trejdosiewicz

Institute: Cancer Research UK, London Research Institute: Lincoln's Inn Fields ; University College London (UCL)

Images:

Tool details

***FOR RESEARCH USE ONLY**

Name: Anti-Basal body marker [LhS 28]

Alternate name:

Class: Monoclonal

Conjugate: Unconjugated

Description: Lhs28 is useful in the identification of ciliated cells in cell associated epithelial pathological specimens.

Purpose:

Parental cell:

Organism:

Tissue:

Model:

Gender:

Isotype: IgG1

Reactivity: Hamster ; Human

Selectivity:

Host: Mouse

Immunogen: BHK a21 Cytoskeletal preparation

Immunogen UNIPROT ID:

Sequence:

Growth properties:

Production details:

Formulation:

Recommended controls:

Bacterial resistance:

Selectable markers:

Additional notes:

Target details

Target: Basal body marker in all ciliated cells.

Target alternate names:

Target background:

Molecular weight:

Ic50:

Applications

Application: IHC ; IHC ; IF ; IP

Application notes:

Handling

Format: Liquid

Concentration: 1 mg/ml

Passage number:

Growth medium:

Temperature:

Atmosphere:

Volume:

Storage medium:

Storage buffer: PBS with 0.02% azide

Storage conditions: -15° C to -25° C

Shipping conditions: Shipping at 4° C

Related tools

Related tools:

References

References: Tao et al. 2013. Cell Res. 23(5):620-34. PMID: 23357851. ; Jiang et al. 2013. Nature. 493(7434):689-93. PMID: 23334421. ; Def defines a conserved nucleolar pathway that leads p53 to proteasome-independent degradation. ; Reciprocal regulation of p53 and malic enzymes modulates

metabolism and senescence. ; Das et al. 2007. Int J Surg Pathol. 15(4):335-45. PMID: 17913939. ; p53 and mdm2 expression in colorectal carcinoma: a correlative analysis with clinical staging and histological parameters. ; Hashemi et al. 2002. Cancer Lett. 180(2):211-21. PMID: 12175554. ; A melanoma-predisposing germline CDKN2A mutation with Fv significance for both p16 and p14ARF. ; Zhang et al. 2001. J Biol Chem. 276(32):29702-10. PMID: 11359766. ; Cyclin a-CDK phosphorylation regulates MDM2 protein interactions. ; Bttger et al. 1997. J Mol Biol. 269(5):744-56. PMID: 9223638. ; Molecular characterization of the hdm2-p53 interaction. ; Dowell et al. 1996. Cytopathology. 7(5):340-51. PMID: 8911758. ; Expression of p21waf1/Cip1, MDM2 and p53 in vivo: analysis of cytological preparations. ; Picksley et al. 1994. Oncogene. 9(9):2523-9. PMID: 8058315. ; Immunochemical analysis of the interaction of p53 with MDM2;--fine mapping of the MDM2 binding site on p53 using synthetic peptides.

CancerTools.org