

# Anti-CD11a (Integrin Subunit Alpha L ) [SPVL7]

**Catalogue number:** 154792

**Sub-type:** Primary antibody

**Images:**

## Contributor

**Inventor:**

**Institute:** Netherlands Cancer Institute

**Images:**

## Tool details

**\*FOR RESEARCH USE ONLY**

**Name:** Anti-CD11a (Integrin Subunit Alpha L ) [SPVL7]

**Alternate name:** ITGAL; LFA-1A

**Class:** Monoclonal

**Conjugate:** Unconjugated

**Description:** ITGAL encodes the integrin alpha L chain. Integrins are heterodimeric integral membrane proteins composed of an alpha chain and a beta chain. This I-domain containing alpha integrin combines with the beta 2 chain (ITGB2) to form the integrin lymphocyte function-associated antigen-1 (LFA-1), which is expressed on all leukocytes. LFA-1 plays a central role in leukocyte intercellular adhesion through interactions with its ligands, ICAMs 1-3 (intercellular adhesion molecules 1 through 3), and also functions in lymphocyte costimulatory signalling.

**Purpose:**

**Parental cell:**

**Organism:**

**Tissue:**

**Model:**

**Gender:**

**Isotype:** IgG1

**Reactivity:** Human

**Selectivity:**

**Host:** Mouse

**Immunogen:** Immunization of BALB/c mice with cells of the T4+T8- cytotoxic T lymphocyte clone HG-38

**Immunogen UNIPROT ID:**

**Sequence:**

**Growth properties:**

**Production details:**

**Formulation:**

**Recommended controls:**

**Bacterial resistance:**

**Selectable markers:**

**Additional notes:**

## Target details

**Target:** CD11a

**Target alternate names:**

**Target background:** ITGAL encodes the integrin alpha L chain. Integrins are heterodimeric integral membrane proteins composed of an alpha chain and a beta chain. This I-domain containing alpha integrin combines with the beta 2 chain (ITGB2) to form the integrin lymphocyte function-associated antigen-1 (LFA-1), which is expressed on all leukocytes. LFA-1 plays a central role in leukocyte intercellular adhesion through interactions with its ligands, ICAMs 1-3 (intercellular adhesion molecules 1 through 3), and also functions in lymphocyte costimulatory signalling.

**Molecular weight:** 126 kDa

**Ic50:**

## Applications

**Application:** FACS ; IP

**Application notes:**

## Handling

**Format:** Liquid

**Concentration:** 0.9-1.1 mg/ml

**Passage number:**

**Growth medium:**

**Temperature:**

**Atmosphere:**

**Volume:**

**Storage medium:**

**Storage buffer:** PBS with 0.02% azide

**Storage conditions:** -15° C to -25° C

**Shipping conditions:** Shipping at 4° C

## Related tools

**Related tools:**

## References

**References:** Lub et al. 1997. Mol Biol Cell. 8(4):719-28. PMID: 9247650.

CancerTools.org