

# Anti-CD3 e [NKIT3a]

**Catalogue number:** 154754

**Sub-type:** Primary antibody

**Images:**

## Contributor

**Inventor:**

**Institute:** Netherlands Cancer Institute

**Images:**

## Tool details

**\*FOR RESEARCH USE ONLY**

**Name:** Anti-CD3 e [NKIT3a]

**Alternate name:** CD3E; T-Cell Surface Glycoprotein CD3 Epsilon Chain

**Class:** Monoclonal

**Conjugate:** Unconjugated

**Description:** The CD3-epsilon polypeptide together with CD3-gamma, -delta and -zeta, and the T-cell receptor alpha/beta and gamma/delta heterodimers, forms the T cell receptor-CD3 complex. This complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways. The genes encoding the epsilon, gamma and delta polypeptides are located in the same cluster on chromosome 11. The epsilon polypeptide plays an essential role in T-cell development. Defects in this gene cause severe immunodeficiency.

**Purpose:**

**Parental cell:**

**Organism:**

**Tissue:**

**Model:**

**Gender:**

**Isotype:** IgG2b

**Reactivity:** Human

**Selectivity:**

**Host:** Mouse

**Immunogen:** Human cytotoxic T-lymphocytes

**Immunogen UNIPROT ID:**

**Sequence:**

**Growth properties:**

**Production details:**

**Formulation:**

**Recommended controls:**

**Bacterial resistance:**

**Selectable markers:**

**Additional notes:**

## Target details

**Target:** CD3 e

**Target alternate names:**

**Target background:** The CD3-epsilon polypeptide together with CD3-gamma, -delta and -zeta, and the T-cell receptor alpha/beta and gamma/delta heterodimers, forms the T cell receptor-CD3 complex. This complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways. The genes encoding the epsilon, gamma and delta polypeptides are located in the same cluster on chromosome 11. The epsilon polypeptide plays an essential role in T-cell development. Defects in this gene cause severe immunodeficiency.

**Molecular weight:**

**Ic50:**

## Applications

**Application:** FACS ; IP

**Application notes:**

## Handling

**Format:** Liquid

**Concentration:** 0.9-1.1 mg/ml

**Passage number:**

**Growth medium:**

**Temperature:**

**Atmosphere:**

**Volume:**

**Storage medium:**

**Storage buffer:** PBS with 0.02% azide

**Storage conditions:** -15° C to -25° C

**Shipping conditions:** Shipping at 4° C

## Related tools

**Related tools:**

## References

**References:** Yssel et al. 1987. Cell Immunol. 105(1):161-73. PMID: 3545502.

CancerTools.org