

Anti-CD51 (Vitronection receptor) [NKIM7]

Catalogue number: 154784

Sub-type: Primary antibody

Images:

Contributor

Inventor:

Institute: Netherlands Cancer Institute

Images:

Tool details

***FOR RESEARCH USE ONLY**

Name: Anti-CD51 (Vitronection receptor) [NKIM7]

Alternate name: ITGAV; Integrin Subunit Alpha V

Class: Monoclonal

Conjugate: Unconjugated

Description: ITGAV encodes integrin alpha chain V. Integrins are heterodimeric integral membrane proteins composed of an alpha chain and a beta chain. Alpha V undergoes post-translational cleavage to yield disulfide-linked heavy and light chains, that combine with multiple integrin beta chains to form different integrins. Among the known associating beta chains (beta chains 1,3,5,6, and 8; 'ITGB1', 'ITGB3', 'ITGB5', 'ITGB6', and 'ITGB8'), each can interact with extracellular matrix ligands; the alpha V beta 3 integrin, perhaps the most studied of these, is referred to as the Vitronection receptor (VNR). In addition to adhesion, many integrins are known to facilitate signal transduction

Purpose:

Parental cell:

Organism:

Tissue:

Model:

Gender:

Isotype: IgG1

Reactivity: Human

Selectivity:

Host: Mouse

Immunogen: Cultured melanoma cells derived from a human melanoma metastasis.

Immunogen UNIPROT ID:

Sequence:

Growth properties:

Production details:

Formulation:

Recommended controls:

Bacterial resistance:

Selectable markers:

Additional notes:

Target details

Target: CD51

Target alternate names:

Target background: ITGAV encodes integrin alpha chain V. Integrins are heterodimeric integral membrane proteins composed of an alpha chain and a beta chain. Alpha V undergoes post-translational cleavage to yield disulfide-linked heavy and light chains, that combine with multiple integrin beta chains to form different integrins. Among the known associating beta chains (beta chains 1,3,5,6, and 8; 'ITGB1', 'ITGB3', 'ITGB5', 'ITGB6', and 'ITGB8'), each can interact with extracellular matrix ligands; the alpha V beta 3 integrin, perhaps the most studied of these, is referred to as the Vitronectin receptor (VNR). In addition to adhesion, many integrins are known to facilitate signal transduction

Molecular weight: 116 kDa

Ic50:

Applications

Application: FACS ; IHC ; IP

Application notes:

Handling

Format: Liquid

Concentration: 0.9-1.1 mg/ml

Passage number:

Growth medium:

Temperature:

Atmosphere:

Volume:

Storage medium:

Storage buffer: PBS with 0.02% azide

Storage conditions: -15° C to -25° C

Shipping conditions: Shipping at 4° C

Related tools

Related tools:

References

References: de Vries et al. 1986. Int J Cancer. 38(4):465-73. PMID: 2428758. ; Buraggi et al. 1985. Cancer Res. 45(7):3378-87. PMID: 4005860.

CancerTools.org