

# Anti-CD56 [NKI-Nbl-3]

**Catalogue number:** 160675

**Sub-type:** Primary antibody

**Images:**

## Contributor

**Inventor:**

**Institute:** Netherlands Cancer Institute

**Images:**

## Tool details

**\*FOR RESEARCH USE ONLY**

**Name:** Anti-CD56 [NKI-Nbl-3]

**Alternate name:** NCAM1; Neural Cell Adhesion Molecule 1

**Class:** Monoclonal

**Conjugate:** Unconjugated

**Description:** CD56, is a homophilic binding glycoprotein expressed on the surface of neurons, glia and skeletal muscle. Although CD56 is often considered a marker of neural lineage commitment due to its discovery site, CD56 expression is also found in, among others, the hematopoietic system. Here, the expression of CD56 is most stringently associated with, but certainly not limited to, natural killer cells. CD56 has been detected on other lymphoid cells, including gamma delta (CD3<sup>+</sup>CD4<sup>+</sup>CD8<sup>+</sup>) T cells and activated CD8<sup>+</sup> T cells, as well as on dendritic cells. CD56 has been implicated as having a role in cell-cell adhesion, neurite outgrowth, synaptic plasticity, and learning and memory

**Purpose:**

**Parental cell:**

**Organism:**

**Tissue:**

**Model:**

**Gender:**

**Isotype:** IgG2

**Reactivity:** Human

**Selectivity:**

**Host:** Mouse

**Immunogen:** CHP-212 human neuroblastoma cell line

**Immunogen UNIPROT ID:**

**Sequence:**

**Growth properties:**

**Production details:**  
**Formulation:**  
**Recommended controls:**  
**Bacterial resistance:**  
**Selectable markers:**  
**Additional notes:**

## Target details

**Target:** CD56

**Target alternate names:**

**Target background:** CD56, is a homophilic binding glycoprotein expressed on the surface of neurons, glia and skeletal muscle. Although CD56 is often considered a marker of neural lineage commitment due to its discovery site, CD56 expression is also found in, among others, the hematopoietic system. Here, the expression of CD56 is most stringently associated with, but certainly not limited to, natural killer cells. CD56 has been detected on other lymphoid cells, including gamma delta ( $\gamma\delta$ ) cells and activated CD8+ T cells, as well as on dendritic cells. CD56 has been implicated as having a role in cellcell adhesion, neurite outgrowth, synaptic plasticity, and learning and memory

**Molecular weight:** ~150 kDa

**Ic50:**

## Applications

**Application:** IP ; WB

**Application notes:**

## Handling

**Format:** Liquid

**Concentration:** 0.9-1.1 mg/ml

**Passage number:**

**Growth medium:**

**Temperature:**

**Atmosphere:**

**Volume:**

**Storage medium:**

**Storage buffer:** PBS with 0.02% azide

**Storage conditions:** -15° C to -25° C

**Shipping conditions:** Shipping at 4° C

## Related tools

Related tools:

## References

References:

CancerTools.org