

# Anti-CD79a [HM57] rAb

**Catalogue number:** 154817

**Sub-type:** Primary antibody

**Images:**

## Contributor

**Inventor:**

**Institute:** Absolute Antibody; University of Oxford

**Images:**

## Tool details

**\*FOR RESEARCH USE ONLY**

**Name:** Anti-CD79a [HM57] rAb

**Alternate name:** CD79a Molecule; Membrane-Bound Immunoglobulin-Associated Protein; Surface IgM-Associated Protein; MB-1 Membrane Glycoprotein; Ig-Alpha; IGA; CD79a Antigen; MB1

**Class:** Recombinant

**Conjugate:** Unconjugated

**Description:** The B-cell Antigen receptor constitutes a disulphide linked heterodimer, consisting of CD79a (mb1) and CD79b / B29 polypeptides which are non-covalently associated with membrane bound immunoglobulins on B-cells. CD79a first appears at pre B-cell stage and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in B-cell lymphomas, in B-cell lines, the majority of acute leukemias of precursor B-cell type and in some myelomas. The CD79a/b heterodimer interacts with at least one tyrosine kinase (Lyn). The induction of tyrosine kinase activity after antigen binding leads to phosphorylation of the CD79a/b dimer, and also of other molecules, thereby initiating intracellular signalling. CD79a is widely used as an adjunct to CD20 as a biomarker for normal and neoplastic B-cells in tissues sections.

**Purpose:**

**Parental cell:**

**Organism:**

**Tissue:**

**Model:**

**Gender:**

**Isotype:** IgG1

**Reactivity:** Bovine ; Chicken ; Horse ; Human ; Guinea Pig ; Mouse ; Opossum ; Pig ; Primate ; Rat ; Rabbit

**Selectivity:**

**Host:** Mouse

**Immunogen:**

Synthetic peptide of 14 amino acids representing residues 202-216 of the human mb-1 cDNA sequence.

**Immunogen UNIPROT ID:**

**Sequence:**

**Growth properties:**

**Production details:**

**Formulation:**

**Recommended controls:**

**Bacterial resistance:**

**Selectable markers:**

**Additional notes:**

## Target details

**Target:** CD79a (mb1)

**Target alternate names:**

**Target background:** The B-cell Antigen receptor constitutes a disulphide linked heterodimer, consisting of CD79a (mb1) and CD79b / B29 polypeptides which are non-covalently associated with membrane bound immunoglobulins on B-cells. CD79a first appears at pre B-cell stage and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in B-cell lymphomas, in B-cell lines, the majority of acute leukemias of precursor B-cell type and in some myelomas. The CD79a/b heterodimer interacts with at least one tyrosine kinase (Lyn). The induction of tyrosine kinase activity after antigen binding leads to phosphorylation of the CD79a/b dimer, and also of other molecules, thereby initiating intracellular signalling. CD79a is widely used as an adjunct to CD20 as a biomarker for normal and neoplastic B-cells in tissues sections.

**Molecular weight:**

**Ic50:**

## Applications

**Application:**

**Application notes:**

## Handling

**Format:** Liquid

**Concentration:**

**Passage number:**

**Growth medium:**

**Temperature:**

**Atmosphere:**

**Volume:**

**Storage medium:**

**Storage buffer:**

**Storage conditions:**

**Shipping conditions:** Shipping at 4° C

## Related tools

**Related tools:**

## References

**References:** Brouns et al. 1993. Eur J Immunol. 23(5):1088-97. PMID: 8477803. ; Evidence for a direct physical interaction of membrane IgM, IgD, and IgG with the B29 gene product. ; Lankester et al. 1994. J Immunol. 152(5):2157-62. PMID: 8133032. ; Mason et al. 1991. J Immunol. 147(11):2474-82. PMID: 1747162. ; The IgM-associated protein mb-1 as a marker of normal and neoplastic B cells. ; The membrane IgM-associated heterodimer on human B cells is a newly defined B cell antigen that contains the protein product of the mb-1 gene. ; The structure of the mu/pseudo light chain complex on human pre-B cells is consistent with a function in signal transduction. ; van Noesel et al. 1991. J Immunol. 146(11):3881-8. PMID: 2033258.