# Anti-Cdc20 [BA8]

Catalogue number: 151302 Sub-type: Primary antibody Images:

# Contributor

Inventor: Julian Gannon Institute: Cancer Research UK, London Research Institute: Clare Hall Laboratories Images:

# **Tool details**

#### **\*FOR RESEARCH USE ONLY**

Name: Anti-Cdc20 [BA8]

ols.org Alternate name: APC/C activator protein CDC2; Cell division control protein 2; YGL116W; Cell Division Cycle 2; P55CDC; CDC2 Cell Division Cycle 2; BA276H19.3; CDC2A

Class: Monoclonal **Conjugate:** Unconjugated Description: Fizzy protein (cdc20) is associated with the activation of the anaphase-promoting complex (APC) at the onset of anaphase. Purpose: Parental cell: Organism: Tissue: Model: Gender: Isotype: IgG2a Reactivity: Xenopus laevis Selectivity: Host: Mouse **Immunogen:** Peptide from the carboxy terminus of the protein. Immunogen UNIPROT ID: Sequence: Growth properties: **Production details:** Formulation: **Recommended controls: Bacterial resistance:** Selectable markers:

#### Additional notes:

#### **Target details**

Target: Cell division cycle protein 20 (cdc20)

Target alternate names:

Target background: Fizzy protein (cdc20) is associated with the activation of the anaphase-promoting complex (APC) at the onset of anaphase.

Molecular weight:

Ic50:

# **Applications**

Application: ELISA ; FACS ; WB **Application notes:** 

## Handling

Cancer Tools.org Format: Liquid **Concentration:** 1 mg/ml Passage number: Growth medium: Temperature: Atmosphere: Volume: Storage medium: Storage buffer: PBS with 0.02% azide Storage conditions: -15° C to -25° C Shipping conditions: Shipping at 4° C

## **Related tools**

Related tools:

## **References**

**References:** Esmaili et al. 2010. Cell Cycle. 9(19):3913-20. PMID: 20935510. ; Wu et al. 2009. NatCell Biol. 11(5):644-51. PMID: 19396163.

