

Anti-ErbB2 [ICR55]

Catalogue number: 153391

Sub-type: Primary antibody

Images:

Contributor

Inventor: Chris Dean

Institute: The Institute of Cancer Research

Images:

Tool details

***FOR RESEARCH USE ONLY**

Name: Anti-ErbB2 [ICR55]

Alternate name: ERBB2, Tyrosine kinase type cell surface receptor HER2, Human Epidermal Growth Factor Receptor 2, Proto-Oncogene C-ErbB-2, HER-2/Neu, HER2, Receptor Tyrosine-Protein Kinase ErbB-2

Class: Monoclonal

Conjugate: Unconjugated

Description: Protein tyrosine kinase that is part of several cell surface receptor complexes, but that apparently needs a coreceptor for ligand binding. Essential component of a neuregulin-receptor complex, although neuregulins do not interact with it alone. GP30 is a potential ligand for this receptor. Regulates outgrowth and stabilization of peripheral microtubules (MTs). Upon ERBB2 activation, the MEMO1-RHOA-DIAPH1 signaling pathway elicits the phosphorylation and thus the inhibition of GSK3B at cell membrane. This prevents the phosphorylation of APC and CLASP2, allowing its association with the cell membrane. In turn, membrane-bound APC allows the localization of MACF1 to the cell membrane, which is required for microtubule capture and stabilization.

Purpose:

Parental cell:

Organism:

Tissue:

Model:

Gender:

Isotype: IgG2a

Reactivity: Human

Selectivity:

Host: Rat

Immunogen: Recombinant full length protein (Human)

Immunogen UNIPROT ID:

Sequence:
Growth properties:
Production details:
Formulation:
Recommended controls:
Bacterial resistance:
Selectable markers:
Additional notes:

Target details

Target: Erb-B2

Target alternate names:

Target background: Protein tyrosine kinase that is part of several cell surface receptor complexes, but that apparently needs a coreceptor for ligand binding. Essential component of a neuregulin-receptor complex, although neuregulins do not interact with it alone. GP30 is a potential ligand for this receptor. Regulates outgrowth and stabilization of peripheral microtubules (MTs). Upon ERBB2 activation, the MEMO1-RHOA-DIAPH1 signaling pathway elicits the phosphorylation and thus the inhibition of GSK3B at cell membrane. This prevents the phosphorylation of APC and CLASP2, allowing its association with the cell membrane. In turn, membrane-bound APC allows the localization of MACF1 to the cell membrane, which is required for microtubule capture and stabilization.

Molecular weight: 77 kDa

Ic50:

Applications

Application: IHC ; IP ; WB

Application notes:

Handling

Format: Liquid

Concentration: 0.9-1.1 mg/ml

Passage number:

Growth medium:

Temperature:

Atmosphere:

Volume:

Storage medium:

Storage buffer: PBS with 0.02% azide

Storage conditions:

-15° C to -25° C

Shipping conditions: Shipping at 4° C

Related tools

Related tools:

References

References: Styles et al. 1990. Int J Cancer. 45(2):320-4. PMID: 1689275. ; Rat monoclonal antibodies to the external domain of the product of the C-erbB-2 proto-oncogene.

CancerTools.org