

Anti-Ig/BCR [L1]

Catalogue number: 154775

Sub-type: Primary antibody

Images:

Contributor

Inventor:

Institute: Netherlands Cancer Institute

Images:

Tool details

***FOR RESEARCH USE ONLY**

Name: Anti-Ig/BCR [L1]

Alternate name:

Class: Monoclonal

Conjugate: Unconjugated

Description: The B-cell receptor (BCR) is composed of immunoglobulin molecules that form a type 1 transmembrane receptor protein usually located on the outer surface of B cells. Through biochemical signalling and by physically acquiring antigens from the immune synapses, the BCR controls the activation of B-cell.

Purpose:

Parental cell:

Organism:

Tissue:

Model:

Gender:

Isotype:

Reactivity: Human

Selectivity:

Host: Mouse

Immunogen: B cell non-Hodgkin's lymphoma

Immunogen UNIPROT ID:

Sequence:

Growth properties:

Production details:

Formulation:

Recommended controls:

Bacterial resistance:

Selectable markers:

Additional notes:

Target details

Target: Ig/BCR

Target alternate names:

Target background: The B-cell receptor (BCR) is composed of immunoglobulin molecules that form a type 1 transmembrane receptor protein usually located on the outer surface of B cells. Through biochemical signalling and by physically acquiring antigens from the immune synapses, the BCR controls the activation of B-cell.

Molecular weight:

Ic50:

Applications

Application: FACS ; IP

Application notes:

Handling

Format: Liquid

Concentration: 0.9-1.1 mg/ml

Passage number:

Growth medium:

Temperature:

Atmosphere:

Volume:

Storage medium:

Storage buffer: PBS with 0.02% azide

Storage conditions: -15° C to -25° C

Shipping conditions: Shipping at 4° C

Related tools

Related tools:

References

References: Stam et al. 1986. J Immunol. 137(7):2299-306. PMID: 3760563.

CancerTools.org