

Anti-Lambda light chain [N10/2] rAb

Catalogue number: 154828

Sub-type: Primary antibody

Images:

Contributor

Inventor:

Institute: Absolute Antibody; University of Oxford

Images:

Tool details

***FOR RESEARCH USE ONLY**

Name: Anti-Lambda light chain [N10/2] rAb

Alternate name:

Class: Recombinant

Conjugate: Unconjugated

Description: All five immunoglobulin classes share the same basic four polypeptide chain structure of two heavy-chains and two light chains. There are five heavy chain types, and two light-chain types (Kappa and Lambda) both having a molecular weight of 22.5kDa. This antibody can be used as a primary or secondary antibody.

Purpose:

Parental cell:

Organism:

Tissue:

Model:

Gender:

Isotype: IgG1

Reactivity: Human ; Saccharomyces cerevisiae

Selectivity:

Host: Mouse

Immunogen: Monoclonal lambda light chains from human urine

Immunogen UNIPROT ID:

Sequence:

Growth properties:

Production details:

Formulation:

Recommended controls:

Bacterial resistance:

Selectable markers:

Additional notes:

Target details

Target: Lambda light chain

Target alternate names:

Target background: All five immunoglobulin classes share the same basic four polypeptide chain structure of two heavy-chains and two light chains. There are five heavy chain types, and two light-chain types (Kappa and Lambda) both having a molecular weight of 22.5kDa. This antibody can be used as a primary or secondary antibody.

Molecular weight:

Ic50:

Applications

Application:

Application notes:

Handling

Format: Liquid

Concentration:

Passage number:

Growth medium:

Temperature:

Atmosphere:

Volume:

Storage medium:

Storage buffer:

Storage conditions:

Shipping conditions: Shipping at 4° C

Related tools

Related tools:

References

References: Brune et al. 2007. Eur Cell Mater. 14:78-90 ; Complex formation of platelet membrane glycoproteins IIb and IIIa with fibrinogen. ; discussion 90-1. PMID: 18085506. ; Gatter et al. 1988. Histopathology. 13(3):257-67. PMID: 3192191. ; Immunological study of in vitro maturation of human megakaryocytes. ; In vitro comparison of human fibroblasts from intact and ruptured ACL for use in tissue engineering. ; Nachman et al. 1982. J Clin Invest. 69(2):263-9. PMID: 6460044. ; The immunohistological detection of platelets, megakaryocytes and thrombi in routinely processed specimens. ; Vinci et al. 1984. Br J Haematol. 56(4):589-605. PMID: 6231944.

CancerTools.org