

Anti-MBP [TVG MBP 3]

Catalogue number: 151169

Sub-type: Primary antibody

Images: https://res.cloudinary.com/ximbio/image/upload/c_fit/2d0d7919-32c9-45f0-b0fe-383121d23a40.jpg

Contributor

Inventor: Lionel Crawford

Institute: University of Cambridge

Images: https://res.cloudinary.com/ximbio/image/upload/c_fit/2d0d7919-32c9-45f0-b0fe-383121d23a40.jpg

Tool details

***FOR RESEARCH USE ONLY**

Name: Anti-MBP [TVG MBP 3]

Alternate name:

Class: Monoclonal

Conjugate: Unconjugated

Description: TVG MBP 3 is useful for detection and isolation of recombinant MBP fusion proteins

Purpose: Marker

Parental cell:

Organism:

Tissue:

Model:

Gender:

Isotype: IgM

Reactivity: Human papilloma virus

Selectivity:

Host: Mouse

Immunogen: A single intravenous injection of the maltose binding protein MBP-E2.

Immunogen UNIPROT ID:

Sequence:

Growth properties:

Production details:

Formulation:

Recommended controls:

Bacterial resistance:

Selectable markers:

Additional notes:

Target details

Target: Maltose binding protein.

Target alternate names:

Target background: TVG MBP 3 is useful for detection and isolation of recombinant MBP fusion proteins

Molecular weight:

Ic50:

Applications

Application: ELISA ; IP ; WB

Application notes:

Handling

Format: Liquid

Concentration: 1 mg/ml

Passage number:

Growth medium:

Temperature:

Atmosphere:

Volume:

Storage medium:

Storage buffer: PBS with 0.02% azide

Storage conditions: -15° C to -25° C

Shipping conditions: Shipping at 4° C

Related tools

Related tools:

References

References: A deletion in the gene encoding the CD45 antigen in a patient with SCID. ; Hibma et al. 1995. Eur J Biochem. 229(2):517-25. PMID: 7744075. ; The interaction between human papillomavirus

type 16 E1 and E2 proteins is blocked by an antibody to the N-terminal region of E2.

CancerTools.org