

Anti-Rad17 [Rad17 1C6/2]

Catalogue number: 151235

Sub-type: Primary antibody

Images:

Contributor

Inventor:

Institute: Cancer Research UK, London Research Institute: Clare Hall Laboratories

Images:

Tool details

***FOR RESEARCH USE ONLY**

Name: Anti-Rad17 [Rad17 1C6/2]

Alternate name:

CancerTools.org

Class: Monoclonal

Conjugate: Unconjugated

Description: RAD17 is a component of the DNA replication and DNA damage checkpoints in yeast.

Purpose:

Parental cell:

Organism:

Tissue:

Model:

Gender:

Isotype: IgG1

Reactivity: Saccharomyces cerevisiae

Selectivity:

Host: Mouse

Immunogen:

Immunogen UNIPROT ID:

Sequence:

Growth properties:

Production details:

Formulation:

Recommended controls:

Bacterial resistance:

Selectable markers:

Additional notes:

Target details

Target: Rad17

Target alternate names:

Target background: RAD17 is a component of the DNA replication and DNA damage checkpoints in yeast.

Molecular weight:

Ic50:

Applications

Application: WB

Application notes:

Handling

Format: Liquid

Concentration:

0.9-1.1 mg/ml

Passage number:

Growth medium:

Temperature:

Atmosphere:

Volume:

Storage medium:

Storage buffer: PBS with 0.02% azide

Storage conditions: -15° C to -25° C

Shipping conditions: Shipping at 4° C

Related tools

Related tools:

References

References: "Monoclonal Antibodies Diagnostics for Individual Members of the Cytochromes P450IV Gene Family", Abstract 1.36 at the 12th Eur. Work. on Drug Metabolism, Basel, Switzerland, Sept. 16-21, 1990. ; "Monoclonal Antibodies Diagnostics for Individual Members of the Cytochromes P450IV Gene Family", Abstract 205 at the VIIIth Inter. Symp. on Microsomes and Drug Oxidations, Stockholm, Sweden, June 25-26, 1990. ; Tzeng et al. 2012. Evid Based Complement Alternat Med. 2012:827278. PMID: 22253647. ; Vinegar-Baked Radix Bupleuri Regulates Lipid Disorders via a Pathway Dependent on Peroxisome-Proliferator-Activated Receptor- α in High-Fat-Diet-Induced Obese Rats.