

N_M5R1 Cell Line

Catalogue number: 152837

Sub-type: Continuous

Images:

Contributor

Inventor: John Lunec

Institute: Newcastle University

Images:

Tool details

***FOR RESEARCH USE ONLY**

Name: N_M5R1 Cell Line

Alternate name:

Class:

Conjugate:

Description: To determine how resistance to MDM2/p53 binding antagonists might develop, NGP cells were exposed to growth inhibitory concentrations of a MDM2 inhibitor, MI-63, and a clonal resistant cell line was generated. The p53 mediated responses of the parental and resistant cell line were compared. In contrast to the parental cell lines, p53 activation by Nutlin-3, MI-63 or ionizing radiation was not observed in the NGP derived cell line, N_M5R1.

Purpose:

Parental cell: NGP

Organism:

Tissue: Lung

Model: Cancer Model

Gender:

Isotype:

Reactivity:

Selectivity:

Host:

Immunogen:

Immunogen UNIPROT ID:

Sequence:

Growth properties:

Production details: Resistant cell lines were established by exposing NGP cells to MI-63. Single cell derived colonies were isolated with cloning cylinders and the clonal population expanded in culture

medium containing the MDM2/p53 antagonist refreshed weekly for 60 days. Stage 1 resistant clones were then further exposed to increased concentrations of MI-63 for 30 days to generate stage 2 resistant clones.

Formulation:

Recommended controls:

Bacterial resistance:

Selectable markers:

Additional notes:

Target details

Target:

Target alternate names:

Target background:

Molecular weight:

Ic50:

Applications

Application:

Application notes:

Handling

Format: Frozen

Concentration:

Passage number:

Growth medium:

Temperature:

Atmosphere:

Volume:

Storage medium:

Storage buffer:

Storage conditions: Liquid Nitrogen

Shipping conditions: Dry ice

Related tools

Related tools:

References

References: Ellis et al. 2012. J Oncol. 2012;65:1507. PMID: 22545050. ; Angiogenesis in Paget's Disease of the Vulva and the Breast: Correlation with Microvessel Density. ; Hussein et al. 2009. Leukemia. 23(5):852-5. PMID: 19194467. ; MPLW515L mutation in acute megakaryoblastic leukaemia. ; Disse et al. 2009. Blood. 113(4):973-80. PMID: 18945966. ; Phospholipase D1 is specifically required for regulated secretion of von Willebrand factor from endothelial cells.

CancerTools.org