

# RMA-S.Trh4 cells

**Catalogue number:** 157675

**Sub-type:** Continuous

**Images:**

## Contributor

**Inventor:** Thorbald Van Hall

**Institute:** Leiden University and Leiden University Medical Center; Stichting Oncode Institute

**Images:**

## Tool details

**\*FOR RESEARCH USE ONLY**

**Name:** RMA-S.Trh4 cells

**Alternate name:**

**Class:**

**Conjugate:**

**Description:** This cell line provides a model to help to understand T-cell recognition through Trh4 expression. ER-resident ceramide synthase Trh4 is a prototypic example of a neo-antigen selectively presented by cells through various pathways. RMA-S lymphoma cells are TAP-deficient on account of an incomplete TAP2 chain, whereas RMA cells have their TAP chain intact. The Trh4 peptide is only presented on cells with impaired TAP function through the classical MHC I pathway. This cell line presents said Trh4...

**Purpose:**

**Parental cell:** RMA-S

**Organism:** Mouse

**Tissue:** Lymphatic Tissue

**Model:** Tumour line

**Gender:**

**Isotype:**

**Reactivity:**

**Selectivity:**

**Host:**

**Immunogen:**

**Immunogen UNIPROT ID:**

**Sequence:**

**Growth properties:**

**Production details:** Retroviral transduction of the mouse Trh4 gene in an IRES-GFP construct

**Formulation:**

**Recommended controls:** RMA.Trh4 cells

**Bacterial resistance:**

**Selectable markers:**

**Additional notes:**

## Target details

**Target:** Trh4

**Target alternate names:**

**Target background:**

**Molecular weight:**

**Ic50:**

## Applications

**Application:**

**Application notes:**

## Handling

**Format:** Frozen

**Concentration:**

**Passage number:**

**Growth medium:**

**Temperature:**

**Atmosphere:**

**Volume:**

**Storage medium:**

**Storage buffer:**

**Storage conditions:** Liquid Nitrogen

**Shipping conditions:** Dry ice

## Related tools

**Related tools:**

## References

**References:** Oliveira et al. 2011. Eur J Immunol. 41(11):3114-24. PMID: 21898382.

CancerTools.org