Sdhb fl/fl Cell Line

Catalogue number: 153294

Sub-type: Images:

Contributor

Inventor: Eyal Gottlieb

Institute: Cancer Research UK, Glasgow: The Beatson Institute

Images:

Tool details

*FOR RESEARCH USE ONLY

Name: Sdhb fl/fl Cell Line

Alternate name: Succinate dehydrogenase, SDH, Succinate-coenzyme Q reductase, SQR

Class:

Conjugate:

Description: Succinate dehydrogenase (SDH) is an enzyme complex found in the inner mitochondrial membrane of mammalian mitochondria and many bacterial cells. SDH is the only enzyme that participates in both the electron transport chain and citric acid cycle. The enzymatic function of SDH oxidises succinate to fumarate in the tricarboxylic acid cycle and loosing this function is associated with cancer formation.

ds.org

Purpose:
Parental cell:
Organism: Mouse
Tissue: Kidney

Model: Conditional KO

Gender: Isotype: Reactivity: Selectivity: Host:

.

Immunogen:

Immunogen UNIPROT ID:

Sequence:

Growth properties: Production details:

Formulation:

Recommended controls:

Bacterial resistance: Selectable markers: Additional notes:

Target details

Target: Succinate dehydrogenase

Target alternate names:

Target background:

Molecular weight:

Ic50:

Applications

Application:

Application notes:

Handling

Format: Frozen
Concentration:
Passage number:
Growth medium:
Temperature:
Atmosphere:
Volume:

Storage medium: Storage buffer:

Storage conditions: Liquid Nitrogen

Shipping conditions: Dry ice

Related tools

Related tools: Sdhb -/- CL 5 Cell Line; Sdhb -/- CL 7 Cell Line; SDHB Floxed Mouse

Cancer Tools.org

References

References: Cardaci et al. 2015. Nat Cell Biol. 17(10):1317-26. PMID: 26302408. ; Pyruvate

carboxylation enables growth of SDH-deficient cells by supporting aspartate biosynthesis.

