# **SDHB Floxed Mouse**

Catalogue number: 153295 Sub-type: Mouse Images:

#### Contributor

Inventor: Eyal Gottlieb Institute: Cancer Research UK, Glasgow: The Beatson Institute Images:

#### **Tool details**

#### **\*FOR RESEARCH USE ONLY**

Name: SDHB Floxed Mouse

Alternate name: Succinate dehydrogenase, SDH, Succinate-coenzyme Q reductase, SQR

Class:

#### Conjugate:

**Description:** Succinate dehydrogenase (SDH) is an enzyme complex found in the inner mitochondrial membrane of mammalian mitochondria and many bacterial cells. SDH is the only enzyme that participates in both the electron transport chain and citric acid cycle. The enzymatic function of SDH oxidises succinate to fumarate in the tricarboxylic acid cycle and loosing this function is associated with cancer formation.

ols.org

Purpose: Parental cell: **Organism: Tissue:** Model: Transgenic Gender: **Isotype: Reactivity:** Selectivity: Host: Immunogen: Immunogen UNIPROT ID: Sequence: Growth properties: Production details: Formulation: **Recommended controls:** 

**Bacterial resistance:** Selectable markers: Additional notes:

### **Target details**

Target: Succinate dehydrogenase

Target alternate names:

Target background:

Molecular weight:

Ic50:

## **Applications**

**Application: Application notes:** 

### Handling

CancerTools.org Format: **Concentration:** Passage number: Growth medium: **Temperature:** Atmosphere: Volume: Storage medium: Storage buffer: Storage conditions: Shipping conditions: Embryo/Spermatoza- Dry Ice

#### **Related tools**

Related tools: Sdhb -/- CL 5 Cell Line ; Sdhb -/- CL 7 Cell Line ; Sdhb fl/fl Cell Line

### References

References: Cardaci et al. 2015. Nat Cell Biol. 17(10):1317-26. PMID: 26302408. ; Pyruvate

carboxylation enables growth of SDH-deficient cells by supporting aspartate biosynthesis.

Cancer Tools.org