

pCA Vector

Catalogue number: 153961

Sub-type: pcDNA3

Images:

Contributor

Inventor: Dr Lawrence Banks

Institute: International Centre For Genetic Engineering And Biotechnology (ICGEB)

Images:

Tool details

***FOR RESEARCH USE ONLY**

Name: pCA Vector

Alternate name:

Class:

Conjugate:

Description: The pCA plasmid was created by inserting HA and FLAG epitopes into the multiple cloning site of pcDNA3

Purpose:

Parental cell:

Organism:

Tissue:

Model:

Gender:

Isotype:

Reactivity:

Selectivity:

Host:

Immunogen:

Immunogen UNIPROT ID:

Sequence:

Growth properties:

Production details:

Formulation:

Recommended controls:

Bacterial resistance:

Selectable markers: Neomycin

Additional notes:

The pCA plasmid is useful for TAP tagging/purification and in vitro expression

Target details

Target: FLAG and HA tags

Target alternate names:

Target background:

Molecular weight:

Ic50:

Applications

Application:

Application notes: The pCA plasmid was created by inserting HA and FLAG epitopes into the multiple cloning site of pcDNA3

Handling

Format:

Concentration:

Passage number:

Growth medium:

Temperature:

Atmosphere:

Volume:

Storage medium:

Storage buffer:

Storage conditions:

Shipping conditions:

Related tools

Related tools:

References

References: Fath et al. 2006. J Biol Chem. 281(19):13612-9. PMID: 16543236. ; Histone deacetylase inhibitors repress the transactivation potential of hypoxia-inducible factors independently of direct acetylation of HIF-alpha.

CancerTools.org